

## WHAT IS CLAIMED IS:

26. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human melanocortin-4 receptor (MC-4R) agonist wherein the binding of the compound to the human MC-4R is characterized by an IC<sub>50</sub> less than 30 nanomolar (nM) and the binding of the compound to the human MC-1R is characterized by an IC<sub>50</sub> greater than 30 nM.
27. The method of Claim 26 wherein the binding of the compound to the human MC-1R is characterized by an IC<sub>50</sub> greater than 100 nM.
28. The method of Claim 26 wherein the binding of the compound to the human MC-1R is characterized by an IC<sub>50</sub> greater than 1000 nM.
29. The method of Claim 26 wherein the binding of the compound to the human MC-1R is characterized by an IC<sub>50</sub> greater than 2100 nM.
30. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the binding of the compound to the human MC-4R is characterized by an IC<sub>50</sub> less than 30 nM and the binding of the compound to the human MC-3R is characterized by an IC<sub>50</sub> greater than 30 nM.
31. The method of Claim 30 wherein the binding of the compound to the human MC-3R is characterized by an IC<sub>50</sub> greater than 100 nM.
32. The method of Claim 30 wherein the binding of the compound to the human MC-3R is characterized by an IC<sub>50</sub> greater than 540 nM.
33. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the binding of the compound to the human MC-4R is characterized by an IC<sub>50</sub> less than

Rule 1.124

TOTT 66406660

30 nM and the binding of the compound to the human MC-5R is characterized by an IC<sub>50</sub> greater than 30 nM.

9  
34. The method of Claim 33 wherein the binding of the compound  
5 to the human MC-5R is characterized by an IC<sub>50</sub> of greater than 100 nM.

10  
35. The method of Claim 33 wherein the binding of the compound  
to the human MC-5R is characterized by an IC<sub>50</sub> greater than 230 nM.

11  
36. The method of Claim 26 wherein the compound is further  
10 characterized by binding to each of the human MC-2R, MC-3R, and MC-5R with an  
IC<sub>50</sub> greater than 30 nM.

12  
37. The method of Claim 27 wherein the compound is further  
15 characterized by binding to each of the human MC-2R, MC-3R, and MC-5R with an  
IC<sub>50</sub> greater than 100 nM.

13  
38. The method of Claim 28 wherein the compound is further  
20 characterized by binding to each of the human MC-2R and MC-3R with an IC<sub>50</sub>  
greater than 540 nM and binding to the MC-5R with an IC<sub>50</sub> greater than 230 nM.

14  
39. The method of Claim 36 wherein the compound is further  
25 characterized by binding to any other human melanocortin receptor with an IC<sub>50</sub>  
greater than 30 nM.

15  
40. The method of Claim 37 wherein the compound is further  
characterized by binding to any other human melanocortin receptor with an IC<sub>50</sub>  
greater than 100 nM.

16  
41. The method of Claim 38 wherein the compound is further  
30 characterized by binding to any other human melanocortin receptor with an IC<sub>50</sub>  
greater than 500 nM.

17  
42. A method of treating sexual dysfunction in a male or female  
35 subject which comprises administering to the subject in need thereof a therapeutically

effective amount of a compound which is a human MC-4R agonist wherein the compound binds to the human MC-4R with a binding affinity at least 10-fold higher than the compound binds to each of the human MC-1R, MC-2R, MC-3R, and MC-5R.

5

<sup>18</sup>  
~~43~~. The method of Claim 42 wherein the compound binds to the human MC-4R with a binding affinity at least 100-fold higher than the compound binds to each of the human MC-1R, MC-2R, MC-3R, and MC-5R.

10

<sup>19</sup>  
~~44~~. The method of Claim 42 wherein the compound binds to the human MC-4R with a binding affinity at least 1000-fold higher than the compound binds to each of the human MC-1R and MC-2R, at least 580-fold higher than the compound binds to the human MC-3R, and at least 250-fold higher than the compound binds to the human MC-5R.

15

<sup>20</sup>  
~~45~~. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the compound binds to the human MC-4R with a binding affinity at least 10-fold higher than the compound binds to any other human melanocortin receptor.

20

<sup>21</sup>  
~~46~~. The method of Claim 45 wherein the compound binds to the human MC-4R with a binding affinity at least 100-fold higher than the compound binds to any other human melanocortin receptor.

25

<sup>22</sup>  
~~47~~. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the functional activity at the MC-4R is characterized by an EC<sub>50</sub> less than 10 nM and the functional activity at the MC-1R is characterized by an EC<sub>50</sub> greater than 10 nM.

30

<sup>23</sup>  
~~48~~. The method of Claim 47 wherein the functional activity of the compound at the MC-1R is characterized by an EC<sub>50</sub> greater than 100 nM.

- 24  
49. The method of Claim 47 wherein the functional activity of the compound at the MC-1R is characterized by an EC<sub>50</sub> greater than 1200 nM.
- 25  
50. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the functional activity at the MC-4R is characterized by an EC<sub>50</sub> less than 10 nM and the functional activity at the MC-3R is characterized by an EC<sub>50</sub> greater than 10 nM.
- 24  
51. The method of Claim 50 wherein the functional activity of the compound at the MC-3R is characterized by an EC<sub>50</sub> greater than 100 nM.
- 23  
52. The method of Claim 50 wherein the functional activity of the compound at the MC-3R is characterized by an EC<sub>50</sub> greater than 1200 nM.
- 28  
53. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the functional activity at the MC-4R is characterized by an EC<sub>50</sub> less than 10 nM and the functional activity at the MC-5R is characterized by an EC<sub>50</sub> greater than 10 nM.
- 29  
54. The method of Claim 53 wherein the functional activity of the compound at the MC-5R is characterized by an EC<sub>50</sub> greater than 100 nM.
- 30  
55. The method of Claim 53 wherein the functional activity of the compound at the MC-5R is characterized by an EC<sub>50</sub> greater than 520 nM.
- 31  
56. The method of Claim 47 wherein the compound is further characterized by having a functional activity at each of the human MC-2R, MC-3R, and MC-5R with an EC<sub>50</sub> greater than 10 nM.
- 32  
57. The method of Claim 48 wherein the compound is further characterized by having a functional activity at each of the human MC-2R, MC-3R, and MC-5R with an EC<sub>50</sub> greater than 100 nM.

- 33  
38. The method of Claim 49 wherein the compound is further characterized by having a functional activity at the human MC-2R and MC-3R with an EC<sub>50</sub> greater than 1200 nM and a functional activity at the human MC-5R with an EC<sub>50</sub> greater than 520 nM.
- 5 34  
35. A method of treating sexual dysfunction in a male or female subject which comprises administering to the subject in need thereof a therapeutically effective amount of a compound which is a human MC-4R agonist wherein the functional activity at the human MC-4R is characterized by an EC<sub>50</sub> at least 10-fold
- 10 lower than the functional activity at each of the human MC-1R, MC-2R, MC-3R, and MC-5R.
- 15 35  
36. The method of Claim 59 wherein the functional activity at the human MC-4R is characterized by an EC<sub>50</sub> at least 100-fold lower than the functional activity at each of the human MC-1R, MC-2R, MC-3R, and MC-5R.
- 20 36  
37. A method for the oral treatment of sexual dysfunction in a male or female subject which comprises the oral administration to the subject in need thereof a therapeutically effective amount of a compound which is an agonist of the human MC-4R.
- 25 37  
38. The method of Claim 61 wherein the compound is a selective agonist of the human MC-4R.
- 38  
39. The method of Claim 61 wherein the sexual dysfunction is erectile dysfunction.

add  
a1